

PRYM VARIETIES ASSOCIATED TO GRAPHS

RUDI SALOMON

ABSTRACT. We present a Prym construction which associates abelian varieties to vertex-transitive strongly regular graphs. As an application we construct Prym-Tyurin varieties of arbitrary exponent ≥ 3 , generalizing a result by Lange, Recillas and Rochas.

1. INTRODUCTION

We describe a Prym construction which associates abelian varieties to certain graphs. More precisely, given the adjacency matrix $A = (a_{ij})_{i,j=1}^d$ of a vertex-transitive strongly regular graph $\mathcal{G}$ along with a covering of curves $p : C \rightarrow \mathbb{P}^1$ of degree d and a labelling $\{x_1, \dots, x_d\}$ of an unramified fiber such that the induced monodromy group of p is represented as a subgroup of the automorphism group of $\mathcal{G}$, we construct a symmetric divisor correspondence D on $C \times C$ which then serves to define complementary subvarieties P_+ and P_- of the Jacobian $J(C)$. The correspondence D is defined in such a way that the point (x_i, x_j) appears in D with multiplicity a_{ij} , analogous to Kanev's construction in [5]. The varieties $P_{\pm}$ are given by $P_{\pm} = \ker(\gamma - r_{\pm} \text{id}_{J(C)})_0$, where $r_{\pm}$ are special eigenvalues of A and γ is the endomorphism on $J(C)$ canonically associated to D (i.e., sending the divisor class $[x - x_0]$ to the class $[D(x) - D(x_0)]$). It is easy to show that

$$(\gamma - r_+ \text{id}_{J(C)})(\gamma - r_- \text{id}_{J(C)}) = 0$$

and $P_{\pm} = \text{im}(\gamma - r_{\pm} \text{id}_{J(C)})$. In particular, if D is fixed point free and $r_+ = 1$, then P_+ is a Prym-Tyurin variety of exponent $1 - r_-$ for C . Given the ramification of p it is not hard to compute the dimension of $P_{\pm}$.

For a thorough definition of D we consider the Galois closure $\pi : X \rightarrow \mathbb{P}^1$ of p and use the induced representation $\text{Gal}(\pi) \rightarrow \text{Aut}(\mathcal{G})$ to construct symmetric correspondences D_+ and D_- on $X \times X$ (much the way M  rindol does in [8]). With C being a quotient curve of X , the correspondence D is derived from $D_{\pm}$ taking quotients and adding $r_{\pm} \Delta_C$, where Δ_C is the diagonal of $C \times C$; see section 4. Given the endomorphisms $\gamma_{\pm}$ on $J(X)$ canonically associated to $D_{\pm}$, we show that $\text{im } \gamma_{\pm}$ and $P_{\pm}$ are isogenous.

The lattice graphs $L_2(n)$, $n \geq 3$ and their complements $\overline{L_2(n)}$ offer important examples. For instance, applying the method to $\overline{L_2(n)}$ and appropriate coverings $C \rightarrow \mathbb{P}^1$ of degree n^2 with branch loci of cardinality $2(l + 2n - 2)$

for $l \geq 1$, we obtain l -dimensional Prym-Tyurin varieties of exponent n for the curves C ; see section 7. We give a characterization of these varieties and show that for $n = 3$ they coincide with the non-trivial Prym-Tyurin varieties of exponent 3 described by Lange, Recillas and Rochas in [6].

Conventions and notations. The ground field is assumed to be the field $\mathbb{C}$ of complex numbers. By a covering of curves we mean a non-constant morphism of irreducible smooth projective curves. The symbol S_n denotes the symmetric group acting on the letters $1, \dots, n$ for $n \in \mathbb{N}$.

2. STRONGLY REGULAR GRAPHS AND MATRICES

We start our discussion with the definition of a strongly regular graph and its adjacency matrix and collect some properties of such graphs. For additional information we refer to [11].

By definition the set of *strongly regular* graphs $\text{SRG}(d, k, \lambda, \mu)$, $k > 0$, consists of the graphs $\mathcal{G}$ with vertex set $\{v_1, \dots, v_d\}$ such that

- a) the set $\Gamma(v_i)$ of vertices adjacent to v_i has exactly k elements and $v_i \notin \Gamma(v_i)$;
- b) for any two adjacent vertices v_i, v_j there are exactly λ vertices adjacent to both v_i and v_j ;
- c) for any two distinct non-adjacent vertices v_i, v_j there are exactly μ vertices adjacent to both v_i and v_j .

Let $A = (a_{ij}) \in \{0, 1\}^{d \times d}$ be the adjacency matrix of such a strongly regular graph $\mathcal{G}$, i.e., $a_{ij} = 1$ if and only if v_i is adjacent to v_j . Then A is symmetric and $(1, \dots, 1) \in \mathbb{R}^d$ is an eigenvector of A with eigenvalue k . The set of eigenvalues of A is $\{k, r_+, r_-\}$ with $r_- < 0 \leq r_+ \leq k$ and

$$(2.1) \quad r_{\pm} = \frac{1}{2} \left[\lambda - \mu \pm \sqrt{(\lambda - \mu)^2 + 4(k - \mu)} \right],$$

implying that

$$(2.2) \quad (A - r_+ I_d)(A - r_- I_d) = \frac{(k - r_+)(k - r_-)}{d} J_d,$$

where J_d is the $d \times d$ matrix whose entries are equal to 1. Given parameters (d, k, λ, μ) such that $(d, k, \lambda, \mu) \neq (4\mu + 1, 2\mu, \mu - 1, \mu)$, one can show that $r_{\pm} \in \mathbb{Z}$ (cf. [11], Theorem 21.1). In fact, if $(d, k, \lambda, \mu) = (4\mu + 1, 2\mu, \mu - 1, \mu)$, then non-integral values of $r_{\pm}$ can occur; for instance, the *Paley* graph $P(5)$ (see [11], Example 21.3) has parameters $(5, 2, 0, 1)$ and eigenvalues $r_{\pm} = -\frac{1}{2} \pm \frac{1}{2}\sqrt{5}$. However, the Paley graph $P(4\mu + 1) \in \text{SRG}(4\mu + 1, 2\mu, \mu - 1, \mu)$, where $4\mu + 1 = p^{2n}$ with p an odd prime and $n \in \mathbb{N}$, has integer eigenvalues $r_{\pm} = \frac{1}{2}(1 \pm p^n)$.

Many strongly regular graphs stem from geometry. The most classical example is offered by the configuration of 27 lines on a cubic surface.


Example 2.1. Given a non-singular cubic surface $X \subset \mathbb{P}^3$, let $\mathcal{L}$ be the intersection graph of the 27 lines that are contained in X . The configuration of the 27 lines is completely described by the 36 *Schäfli double-sixes*, i.e., pairs $M := (\{a_1, \dots, a_6\}, \{b_1, \dots, b_6\})$ of sets of 6 skew lines on X such that each line from one set is skew to a unique line from the other set. Fix a double-six M ; in matrix notation we may write

$$M = \begin{pmatrix} a_1 & a_2 & a_3 & a_4 & a_5 & a_6 \\ b_1 & b_2 & b_3 & b_4 & b_5 & b_6 \end{pmatrix}$$

such that two lines meet if and only they are in different rows and columns. The remaining 15 lines on X are the $c_{ij} := a_i b_j \cap a_j b_i$ with $i \neq j$, where $a_i b_j$ is the plane spanned by the lines a_i and b_j . In this notation the 36 double-sixes are M , the 15 $M_{i,j}$'s and the 20 $M_{i,j,k}$'s, where the double-sixes $M_{i,j}$ and $M_{i,j,k}$ are respectively given by

$$\begin{pmatrix} a_i & b_i & c_{jk} & c_{jl} & c_{jm} & c_{jn} \\ a_j & b_j & c_{ik} & c_{il} & c_{im} & c_{in} \end{pmatrix}, \quad \begin{pmatrix} a_i & a_j & a_k & c_{mn} & c_{ln} & c_{ln} \\ c_{jk} & c_{ik} & c_{ij} & b_l & b_m & b_n \end{pmatrix}.$$

It follows that the *Schläfli* graph $\mathcal{L}$ is in $\text{SRG}(27, 10, 1, 5)$ (the unique member of this set). It is easily seen that the stabilizer of a double-six is a subgroup of $\text{Aut}(\mathcal{L})$ of index 36, isomorphic to $S_6 \times \mathbb{Z}/2\mathbb{Z}$; consequently $\#(\text{Aut}(\mathcal{L})) = 6! \cdot 2 \cdot 36$. In fact, $\text{Aut}(\mathcal{L})$ is isomorphic to the Weyl group $W(E_6)$: Consider the Dynkin diagram for E_6



By definition $W(E_6)$ is generated by the reflections $s_1, \dots, s_5, s$ associated to the simple roots $x_1, \dots, x_5, y$. One shows that there is a surjective homomorphism $W(E_6) \rightarrow \text{Aut}(\mathcal{L})$ sending s_i (resp. s) to the transformation that interchanges the rows of the double-six $M_{i,i+1}$ (resp. $M_{1,2,3}$) (cf. [7], sections 25,26). Then recall that $W(E_6)$ is of order $6! \cdot 2 \cdot 36$. Under this apparent isomorphism the 27 lines on X correspond to the 27 fundamental weights of E_6 . Since $W(E_6)$ acts transitively on these weights, we may consider $\text{Aut}(\mathcal{L})$ as a transitive subgroup of S_{27} .

Additional properties (of strongly regular graphs):

- 1) if $\mathcal{G}$ is disconnected, then $\mathcal{G}$ is the disjoint union of $m > 1$ copies of the *complete* graph K_{k+1} with adjacency matrix $J_{k+1} - I_{k+1}$, where I_{k+1} is the identity matrix. So $(d, k, \lambda, \mu) = (m(k+1), k, k-1, 0)$ and A has exactly two distinct eigenvalues: $k (= r_+)$ with multiplicity m and $r_- = -1$ with multiplicity $d - m$;
- 2) if $\mathcal{G}$ is connected and $\mathcal{G} \neq K_{k+1}$, then $k \neq r_{\pm}$ and k is a simple eigenvalue;
- 3) $\mathcal{G}$ and its complement $\overline{\mathcal{G}} \in \text{SRG}(d, d-k-1, d-2k+\mu-2, d-2k+\lambda)$ are connected if and only if $0 < \mu < k < d-1$, in which case $\mathcal{G}$ is said to be *non-trivial*.

Let $\text{Aut}(A)$ be the stabilizer of A under the natural operation of S_d on $d \times d$ integer matrices by $(a_{ij}) \mapsto (a_{\sigma(i)\sigma(j)})$ and observe that $\text{Aut}(A)$ coincides with $\text{Aut}(\mathcal{G})$. For disconnected $\mathcal{G}$ it is easily seen that $\text{Aut}(A)$ is a transitive subgroup of S_d , implying that disconnected strongly regular graphs are *vertex-transitive*. In practice it turns out that vertex-transitivity is quite rare among non-trivial strongly regular graphs, although most of the sets $\text{SRG}(d, k, \lambda, \mu)$ of non-trivial strongly regular graphs have a vertex-transitive member.

Definition 2.2. Let $A \in \mathbb{Z}^{d \times d}$ be a symmetric matrix with transitive stabilizer group $\text{Aut}(A)$. Then A is a *Prym matrix* if there exist integers k, r_+, r_- with $r_+ > r_-$ such that equation (2.2) holds. Further, if A is a Prym matrix and $m \in \mathbb{N}$, then $A^{\oplus m} := \bigoplus_{i=1}^m A$ is a *repeated Prym matrix*.

Remarks. Suppose that $A \in \mathbb{Z}^{d \times d}$ is a symmetric matrix for which there exist integers k, r_+, r_- with $r_+ > r_-$ such that equation (2.2) holds. Decomposing $\mathbb{R}^d$ into eigenspaces of A we may assume that A takes diagonal form. Then J_d simultaneously transforms into the diagonal matrix $\text{diag}(d, 0, \dots, 0)$, implying that $(1, \dots, 1) \in \mathbb{R}^d$ is an eigenvector of A with, say, eigenvalue η . Hence A has eigenvalues η, r_+, r_- and by (2.2) we have $(\eta - r_+)(\eta - r_-) = (k - r_+)(k - r_-)$, that is, $\eta = k$ or $\eta = r_+ + r_- - k$. Further, if $\eta \neq r_{\pm}$, then η is simple. A $d \times d$ Prym matrix A therefore has integer eigenvalues k, r_+, r_- with $r_+ > r_-$ such that (2.2) holds and k belongs to the eigenvector $(1, \dots, 1)$ of A . Clearly, if A is such a matrix, then for any $m \in \mathbb{N}$ the repeated Prym matrix $A^{\oplus m}$ has the same eigenvalues k, r_+, r_- and satisfies the equation

$$(2.3) \quad (A^{\oplus m} - r_+ I_{md})(A^{\oplus m} - r_- I_{md}) = \frac{(k - r_+)(k - r_-)}{d} J_d^{\oplus m}.$$

Moreover, it is immediately seen that the repeated Prym matrix $A^{\oplus m}$ has transitive stabilizer $\text{Aut}(A^{\oplus m})$. Throughout the paper the eigenvalues of a Prym matrix A will be denoted by k, r_+, r_- , where k belongs to the eigenvector $(1, \dots, 1)$ of A and $r_+ > r_-$.

Many of the known constructions for Prym varieties rely on the definition of a reduced symmetric divisor correspondence for a curve. These correspondences can often be related to Prym matrices whose entries are in the set $\{0, 1\}$. Hence it is useful to have a characterization for such matrices:

Proposition 2.3. Assume that $A = (a_{ij}) \in \{0, 1\}^{d \times d}$ is a Prym matrix with $a_{11} = 0$. Then A is the adjacency matrix of a graph $\mathcal{G} \in \text{SRG}(d, k, \lambda, \mu)$ with $\lambda = k + r_+ r_- + r_+ + r_-$ and $\mu = k + r_+ r_-$.

Proof. By transitivity we have $a_{ii} = 0$ for all $i = 1, \dots, d$, hence A is the adjacency matrix of a regular graph $\mathcal{G}$ of degree k with d vertices. If indeed we have $\mathcal{G} \in \text{SRG}(d, k, \lambda, \mu)$ for some integers λ and μ , then solving equation (2.1) for λ, μ yields $\lambda = k + r_+ r_- + r_+ + r_-$ and $\mu = k + r_+ r_-$. To show that $\mathcal{G}$

is strongly regular we may assume by transitivity that $\mathcal{G}$ is the disjoint union of finitely many copies of a connected graph $\mathcal{G}_c$ with adjacency matrix A_c . If $k \neq r_{\pm}$, then k is simple, so $A = A_c$ and it follows that $\mathcal{G}$ is strongly regular (cf. [11], the remark at the bottom of p. 265). Hence assume that A has just two distinct eigenvalues, i.e., $k = r_{+}$. The complementary graph $\overline{\mathcal{G}}$ has adjacency matrix $J_d - I_d - A$ which is Prym and has eigenvalues $k' = d - k - 1$, $r'_+ = -r_- - 1$ and $r'_- = -k - 1$. So if $r_- \neq k - d$, then $k' \neq r'_{\pm}$, that is, $\overline{\mathcal{G}}$ is a non-complete connected strongly regular graph and so $\mathcal{G}_c = K_{k+1}$. Finally, suppose that $r_- = k - d$. By equation (2.2) we have $A^2 = (2k - d)A - k(k - d)I_d$, hence if v_h, v_i, v_j are three distinct vertices of $\mathcal{G}$ such that v_h is adjacent to v_i and v_i is adjacent to v_j , then the relation $1 = a_{hi} \leq \sum_{l=1}^d a_{hl}a_{lj} = (2k - d)a_{hj}$ implies that v_h is adjacent to v_j . Consequently $\mathcal{G} = K_{k+1}$. $\square$

Example 2.4. Among the set of strongly regular graphs there are some distinguished families of non-trivial vertex transitive graphs. One such family is that of *lattice* graphs; for $n \geq 3$, the lattice graph $L_2(n)$ is the graph with vertex set $\{1, \dots, n\}^2$ such that two distinct vertices (i, j) and (l, m) are adjacent if and only if $i = l$ or $j = m$. Clearly, $S_n \times S_n$ is a transitive subgroup of $\text{Aut}(L_2(n))$, hence the adjacency matrix of $L_2(n)$ and that of its complement $\overline{L_2(n)}$ are Prym. Another subgroup of $\text{Aut}(L_2(n))$ is S_2 ; it permutes the coordinates of the vertices of $L_2(n)$. In fact, it is well-known that the automorphism group of $L_2(n) \in \text{SRG}(n^2, 2(n-1), n-2, 2)$ is equal to the semi-direct product $S_2 \rtimes (S_n \times S_n)$. With reference to future examples (e.g., Examples 5.2 and 5.3) we shall characterize those $\varphi \in \text{Aut}(L_2(n))$ for which each vertex (i, j) of $\overline{L_2(n)}$ (resp. $L_2(n)$ for n odd) is non-adjacent to $\varphi(i, j)$. It is easy to check that:

- i) Each vertex (i, j) of $\overline{L_2(n)}$ is non-adjacent to $\varphi(i, j)$ if and only if $\varphi = (\sigma, \tau)$ with $\sigma, \tau \in S_n$ and $\sigma = (1)$ or $\tau = (1)$.
- ii) Assume that n is odd and φ is a reflection. Then each vertex (i, j) of $L_2(n)$ is non-adjacent to $\varphi(i, j)$ if and only if $\varphi = (\sigma, \sigma^{-1}) \circ t$ with $\sigma \in S_n$ and $t = \begin{pmatrix} 1 & 2 \end{pmatrix} \in S_2$.

Note moreover that $L_2(3)$ and $\overline{L_2(3)}$ are isomorphic; if we identify the set $\{1, \dots, n\}$ with the group $\mathbb{Z}/3\mathbb{Z}$, then the matrix $\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ defines an isomorphism of graphs $L_2(3) \xrightarrow{\sim} \overline{L_2(3)}$.

Example 2.5. Another non-trivial family is that of *Latin square* graphs; for $n \geq 3$, the Latin square graph $L_3(n) \in \text{SRG}(n^2, 3(n-1), n, 6)$ is the graph with vertex set $(\mathbb{Z}/n\mathbb{Z})^2$ such that two distinct vertices (i, j) and (l, m) are adjacent if and only if $i = l$ or $j = m$ or $i + j = l + m$. We identify three subgroups of $\text{Aut}(L_3(n))$; to begin with, the diagonal action of $\text{Aut}(\mathbb{Z}/n\mathbb{Z})$ on $(\mathbb{Z}/n\mathbb{Z})^2$ induces an embedding $\text{Aut}(\mathbb{Z}/n\mathbb{Z}) \hookrightarrow \text{Aut}(L_3(n))$. Similarly, $(\mathbb{Z}/n\mathbb{Z})^2$ induces a subgroup of $\text{Aut}(L_3(n))$ by translation, thus implying that $L_3(n)$ is vertex-transitive. Finally, the subgroup $S := \langle s, t \rangle \subset$

$\text{Aut}((\mathbb{Z}/n\mathbb{Z})^2)$ with s and t respectively given by the matrices $\begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, is immediately seen to be a subgroup of $\text{Aut}(L_3(n))$. Note that sending $s \mapsto (1 \ 2)$ and $t \mapsto (1 \ 3)$ induces an isomorphism $S \xrightarrow{\sim} S_3$. Clearly, the actions of $\text{Aut}(\mathbb{Z}_n)$ and S commute. It can be shown that $\text{Aut}(L_3(n))$ coincides with the semi-direct product $(\mathbb{Z}/n\mathbb{Z})^2 \rtimes (S \times \text{Aut}(\mathbb{Z}/n\mathbb{Z}))$.

3. PRYM VARIETIES OF A GALOIS COVERING

Given a Prym matrix A , we describe a method that associates certain abelian varieties to a finite Galois coverings of $\mathbb{P}^1$ whose Galois group is represented as a transitive subgroup of $\text{Aut}(A^{\oplus m})$, $m \geq 1$. We assume from now on that A is a $d \times d$ Prym matrix with eigenvalues k, r_+, r_- .

Definition 3.1. Consider $\mathcal{P} = (A^{\oplus m}, r, \pi, \phi)$ for $m \geq 1$, where $r \in \{r_+, r_-\}$, π is a finite Galois covering of $\mathbb{P}^1$ and $\phi : \text{Gal}(\pi) \rightarrow S_{md}$ is a transitive representation. Then $\mathcal{P}$ is said to represent *Prym data* if $\phi(G)$ is a subgroup of $\text{Aut}(A^{\oplus m})$.

For instance, assume that we have a finite subset $B = \{b_1, \dots, b_n\}$ of $\mathbb{P}^1$ along with non-trivial permutations $\sigma_1, \dots, \sigma_n \in \text{Aut}(A^{\oplus m})$ such that $\sigma_1 \dots \sigma_n = (1)$ and $G := \langle \sigma_1, \dots, \sigma_n \rangle$ is a transitive subgroup of S_{md} . Let Σ_i be the conjugation class of σ_i in G . According to Riemann's Existence Theorem (RET), the number of equivalence classes of Galois coverings of $\mathbb{P}^1$ of ramification type $\mathcal{R} := [G, B, \{\Sigma_i\}_{i=1}^n]$ is equal to the number of sets $\{(g\tau_1g^{-1}, \dots, g\tau_ng^{-1}) \mid g \in G\}$ with $\tau_i \in \Sigma_i$ such that $\tau_1 \dots \tau_n$ is trivial (cf. [12], p. 37). Hence, let π be a Galois covering of type $\mathcal{R}$. Clearly, if $\phi : \text{Gal}(\pi) \xrightarrow{\sim} G$ is a group isomorphism, then $(A^{\oplus m}, r, \pi, \phi)$ represents Prym data. In this way, varying the continuous parameters $b_1, \dots, b_n$, we obtain finitely many smooth n -dimensional families of Galois coverings with associated Prym data.

The construction. Let $\mathcal{P} = (A^{\oplus m}, r, \pi, \phi)$ be Prym data for a given Galois covering $\pi : X \rightarrow \mathbb{P}^1$. We are going to define two symmetric divisor correspondences, one for X and one for the quotient curve $C := X/H$, where $H \subset G := \text{Gal}(\pi)$ is the stabilizer of the letter 1 with respect to ϕ . Let $(\ , \)_r : \mathbb{R}^{md} \times \mathbb{R}^{md} \rightarrow \mathbb{Q}$ be the symmetric bilinear form canonically associated to the matrix $A^{\oplus m} - rI_{md}$. Given the standard basis $e_1, \dots, e_{md}$ of $\mathbb{R}^{md}$, consider the permutation representation of G on $\mathbb{R}^{md}$ induced by $g : e_i \mapsto e_{g(i)}$. For $g \in G$, denote $\hat{g} = HgH$. Then $(\ , \)_r$ is immediately seen to be G -invariant and $(g_1e_1, e_1)_r = (g_2e_1, e_1)_r$ for all $g_1, g_2 \in G$ such that $\hat{g}_1 = \hat{g}_2$. Let $\alpha : X \rightarrow C$ and $p : C \rightarrow \mathbb{P}^1$ be the canonical mappings. For each $g \in G$ take the graph $\Gamma_g = (\text{id}_X, g)(X)$ of g and let $\hat{\Gamma}_g = (\alpha \times \alpha)(\Gamma_g)$ be its reduced image in $C \times C$. Assume that B is the branch locus of π and put $C_0 = p^{-1}(\mathbb{P}^1 \setminus B)$. Observe that

$$\hat{\Gamma}_{g_1} \cap \hat{\Gamma}_{g_2} \cap (C_0 \times C_0) \neq \emptyset \iff \hat{g}_1 = \hat{g}_2 \iff \hat{\Gamma}_{g_1} = \hat{\Gamma}_{g_2},$$

for all $g_1, g_2 \in G$. Hence we have divisor correspondences $D_{\mathcal{P}}$ on $X \times X$ and $\hat{D}_{\mathcal{P}}$ on $C \times C$, given by

$$D_{\mathcal{P}} = \sum_{g \in G} (ge_1, e_1)_r \Gamma_g, \quad \hat{D}_{\mathcal{P}} = \sum_{\hat{g} \in \hat{G}} (ge_1, e_1)_r \hat{\Gamma}_g$$

with $\hat{G} = H \backslash G / H$. Note that $D_{\mathcal{P}}$ and $\hat{D}_{\mathcal{P}}$ are symmetric as $(ge_1, e_1)_r = (g^{-1}e_1, e_1)_r$ for all $g \in G$.

Definition 3.2. Let $\gamma_{\mathcal{P}}$ on $J(X)$ (resp. $\hat{\gamma}_{\mathcal{P}}$ on $J(C)$) be the endomorphism canonically associated to $D_{\mathcal{P}}$ (resp. $\hat{D}_{\mathcal{P}}$). Then we call $Z_{\mathcal{P}} = \text{im } \gamma_{\mathcal{P}}$ (resp. $\hat{Z}_{\mathcal{P}} = \text{im } \hat{\gamma}_{\mathcal{P}}$) the Prym variety of X (resp. C) associated to $\mathcal{P}$.

Remark. Prym data can be seen as an ‘ornamented’ covering (i.e., a covering with additional data), where the ornamentation is such that we can define divisorial correspondences and Prym varieties.

For $q \in \mathbb{P}^1 \setminus B$ an identification $\pi^{-1}(q) \leftrightarrow G$ is called a *Galois labelling* of the fiber $\pi^{-1}(q)$ if the action of G on the fiber is compatible with its action on itself via multiplication on the left. Moreover, a Galois labelling of $\pi^{-1}(q)$ induces a Galois labelling $p^{-1}(q) \leftrightarrow H \backslash G$ of the fiber of $p : C \rightarrow \mathbb{P}^1$ over q . Denoting $\bar{g} = Hg$ for $g \in G$, we have:

Lemma 3.3. *Given $q \in \mathbb{P}^1 \setminus B$, take a Galois labelling $p^{-1}(q) \leftrightarrow H \backslash G$. Then, for any $\sigma \in G$, the restriction of $\hat{D}_{\mathcal{P}}$ to $\{\bar{\sigma}\} \times C$ is given by the identity*

$$\hat{D}_{\mathcal{P}}(\bar{\sigma}) = \sum_{\bar{g} \in H \backslash G} (ge_1, e_1)_r \bar{g}\bar{\sigma}.$$

Proof. Because $(\bar{\sigma}, \bar{g}\bar{\sigma}) \in \hat{\Gamma}_f \iff \hat{f} = \hat{g}$, for all $f, g \in G$, the point $(\bar{\sigma}, \bar{g}\bar{\sigma})$ appears in $\hat{D}_{\mathcal{P}}$ with multiplicity $(ge_1, e_1)_r$. $\square$

Since H is the stabilizer of a point, the transitivity of the representation $\phi : G \rightarrow S_{md}$ implies that there exists a bijection $H \backslash G \rightarrow \{1, \dots, md\}$. Hence the Galois labelling $p^{-1}(q) \leftrightarrow H \backslash G$ induces a labelling $\{y_1, \dots, y_{md}\}$ of $p^{-1}(q)$ such that $\bar{g}$ corresponds to $y_{g^{-1}(1)}$. The formula in the preceding lemma now turns into

$$(3.1) \quad \hat{D}_{\mathcal{P}}(y_i) = \sum_{j=1}^{md} (e_i, e_j)_r y_j = -ry_i + \sum_{j=1}^{md} ({}^t e_j A^{\oplus m} e_i) y_j,$$

for all $i = 1, \dots, md$.

Proposition 3.4. *Let $\mathcal{P} = (A^{\oplus m}, r, \pi, \phi)$ be Prym data with $\pi : X \rightarrow \mathbb{P}^1$. Then the Prym varieties $Z_{\mathcal{P}}$ of X and $\hat{Z}_{\mathcal{P}}$ of $C := X/H$ are isogenous. Using the notation $\mathcal{P}' = (J_d^{\oplus m}, 0, \pi, \phi)$ and $s = r_+ + r_-$, we have the following quadratic equations for the endomorphisms $\hat{\gamma}_{\mathcal{P}}$ on $J(C)$ and $\gamma_{\mathcal{P}}$ on $J(X)$:*

$$\hat{\gamma}_{\mathcal{P}}(\hat{\gamma}_{\mathcal{P}} + (2r - s)\text{id}_{J(C)}) = \frac{(k - r_+)(k - r_-)}{d} \hat{\gamma}_{\mathcal{P}'}$$

and

$$\gamma_p(\gamma_p + \#(H)(2r - s)\text{id}_{J(X)}) = \#(H) \frac{(k - r_+)(k - r_-)}{d} \gamma_{p'}.$$

Proof. Write $\alpha^* : J(C) \rightarrow J(X)$ for the map induced by α and write $N_\alpha : J(X) \rightarrow J(C)$ for the norm map. Abusing notation, we define homomorphisms $\alpha^* : \mathbb{Z}[H \setminus G] \rightarrow \mathbb{Z}[G]$ and $N_\alpha : \mathbb{Z}[G] \rightarrow \mathbb{Z}[H \setminus G]$, respectively induced by $\bar{g} \mapsto \sum_{h \in H} hg$ and $g \mapsto \bar{g}$. As a direct consequence of Lemma 3.3 we have $N_\alpha D_p = \#(H) \hat{D}_p N_\alpha$. Hence $N_\alpha \gamma_p = \#(H) \hat{\gamma}_p N_\alpha$ and $\gamma_p = \alpha^* \hat{\gamma}_p N_\alpha$. The first identity implies that the restricted mapping $N_\alpha : Z_p \rightarrow \hat{Z}_p$ is surjective, while the two identities combined imply that $\dim \text{im } Z_p = \dim \text{im } \hat{Z}_p$. Therefore the restriction of N_α to Z_p is an isogeny.

Applying equation (3.1) to $\mathcal{P}'$ we obtain $\hat{D}_{p'}(y_i) = \sum_{j=1}^{md} ({}^t e_j J_d^{\oplus m} e_i) y_j$, for all $i = 1, \dots, md$. Hence equations (2.3) and (3.1) imply

$$\hat{D}_p(\hat{D}_p(y)) + (2r - s)\hat{D}_p(y) = \frac{(k - r_+)(k - r_-)}{d} \hat{D}_{p'}(y)$$

for all y in a fiber of $p : C \rightarrow \mathbb{P}^1$ over a point outside the branch locus of π . Let $\hat{\delta}$ (resp. δ) denote the difference between the expressions on the left and the righthand side of the first (resp. second) identity of the proposition. We already know that $\hat{\delta} = 0$. Hence, as $\gamma_{p'} = \alpha^* \hat{\gamma}_{p'} N_\alpha$, we have $\delta = \#(H) \alpha^* \hat{\delta} N_\alpha = 0$. $\square$

Proposition 3.5. *In the special case $m = 1$ the endomorphisms $\hat{\gamma}_{p'}$ and $\gamma_{p'}$ vanish and, using the notation $\mathcal{P}^\pm = (A, r_\pm, \pi, \phi)$, we find that $\hat{Z}_{p+}$ and $\hat{Z}_{p-}$ are complementary subvarieties of $J(C)$. Moreover, $\hat{Z}_{p^\pm} = \ker(\hat{\gamma}_{p^\mp})_0$ and $Z_{p^\pm} = \ker(\gamma_{p^\pm} \pm \#(H)(r_+ - r_-)\text{id}_{J(X)})_0$.*

Proof. Let $q \in \mathbb{P}^1 \setminus B$. By definition of $D_{p'}$, we have $D_{p'}(x) = \pi^* \pi(x)$ for all $x \in \pi^{-1}(q)$, while Lemma 3.3 implies that $\hat{D}_{p'}(y) = p^* p(y)$ for all $y \in p^{-1}(q)$. Hence $\gamma_{p'}$ and $\hat{\gamma}_{p'}$ vanish. As $\hat{\gamma}_{p-} = \hat{\gamma}_{p+} + (r_+ - r_-)\text{id}_{J(C)}$, we note that $\hat{Z}_{p-} = \text{im}(\hat{\gamma}_{p+} + (r_+ - r_-)\text{id}_{J(C)})$. Using standard arguments one shows that $\varepsilon_+ := \frac{-1}{r_+ - r_-} \hat{\gamma}_{p+}$ and $\varepsilon_- := \frac{1}{r_+ - r_-} (\hat{\gamma}_{p+} + (r_+ - r_-)\text{id}_{J(C)})$ are symmetric idempotents in $\text{End}_{\mathbb{Q}}(J(C))$. Since $\varepsilon_+ = \text{id}_{J(C)} - \varepsilon_-$, it follows that Z_{p+} and Z_{p-} are complementary subvarieties of $J(C)$ and $\dim Z_{p+} + \dim Z_{p-} = g(C)$. As a consequence of Proposition 3.4 we have $\hat{Z}_{p^\pm} \subset \ker(\hat{\gamma}_{p^\mp})_0$. Moreover, by [3], Proposition 5.1.1 the analytic representation $\varrho_a(\hat{\gamma}_{p+}) \in \text{End}(H^0(C, \omega_C))$ of $\hat{\gamma}_{p+}$ is self-adjoint with eigenvalues $r_- - r_+$ and 0 with respect to the Riemann form $c_1(\Theta_C)$, where Θ_C is the canonical polarization of $J(C)$. Since $\dim \ker(\hat{\gamma}_{p^\pm})_0 = \dim \ker \varrho_a(\hat{\gamma}_{p^\pm})$, we have $\dim \ker(\hat{\gamma}_{p+})_0 + \dim \ker(\hat{\gamma}_{p-})_0 = g(C)$ and therefore $\hat{Z}_{p^\pm} = \ker(\hat{\gamma}_{p^\mp})_0$. Similarly, the remaining assertion follows from the fact that $Z_{p^\pm}$ and $\text{im}(\gamma_{p^\pm} \pm \#(H)(r_+ - r_-)\text{id}_{J(X)})$ are complementary subvarieties of $J(X)$. $\square$

Finally, we note that the Prym data $\mathcal{P} = (A, r, \pi, \phi)$ associated to the adjacency matrix A of a strongly regular graph $\mathcal{G}$ and the Prym data $\overline{\mathcal{P}} = (J_d - I_d - A, -r - 1, \pi, \phi)$ associated to the adjacency matrix $J_d - I_d - A$ of the complementary graph $\overline{\mathcal{G}}$ yield the same Prym varieties because of the identities $\gamma_{\mathcal{P}} + \gamma_{\overline{\mathcal{P}}} = 0$ and $\hat{\gamma}_{\mathcal{P}} + \hat{\gamma}_{\overline{\mathcal{P}}} = 0$.

4. PRYM VARIETIES OF A NON-GALOIS COVERING

We now shift our attention to non-Galois coverings of $\mathbb{P}^1$. Given a (repeated) Prym matrix $A^{\oplus m}$ and a special covering $p : C \rightarrow \mathbb{P}^1$ of degree md along with an appropriate *labelling class* (to be defined below), we shall define a symmetric divisor correspondence on $C \times C$ which then serves to obtain a pair of Prym varieties in $J(C)$.

To define labelling classes, let $p : C \rightarrow \mathbb{P}^1$ be a covering of degree n with branch locus B . Given a point $q \in \mathbb{P}^1 \setminus B$, a labelling $\{x_1, \dots, x_n\}$ of the fiber $p^{-1}(q)$ induces a bijection $\nu : p^{-1}(q) \rightarrow \{1, \dots, n\}$ sending x_i to i . For $q_j \in \mathbb{P}^1 \setminus B$ with $j = 1, 2$, let $\{x_{j1}, \dots, x_{jn}\}$ be a labelling of the fiber above q_j inducing a bijection ν_j . Then the bijections ν_1 and ν_2 are said to be equivalent if there exists a path $\mu \subset \mathbb{P}^1 \setminus B$ running from q_1 to q_2 such that the lift of μ to C with initial point x_{1i} has end point x_{2i} , for $i = 1, \dots, n$. The equivalence class $[\nu]$ of an induced bijection ν is called a labelling class for p .

Definition 4.1. Consider the triple $\mathcal{T} = (A^{\oplus m}, p, [\nu])$ for $m \geq 1$, where $p : C \rightarrow \mathbb{P}^1$ is a covering of degree md and $[\nu]$ is a labelling class for p . We say that $\mathcal{T}$ is a *Prym triple* if $\text{Aut}(A^{\oplus m})$ contains the monodromy group of p with respect to $[\nu]$.

Let $B = \{b_1, \dots, b_n\}$ be a finite subset of $\mathbb{P}^1$ and take non-trivial permutations $\sigma_1, \dots, \sigma_n \in \text{Aut}(A^{\oplus m})$ such that $\sigma_n \dots \sigma_1 = (1)$ and $G = \langle \sigma_1, \dots, \sigma_n \rangle$ is a transitive subgroup of S_{md} . Recalling the monodromy version of RET (cf. [9], p. 92), we may assume that $p : C \rightarrow \mathbb{P}^1$ is a covering of degree md with branch locus B and labelling class $[\nu]$ such that the ramification of p above b_i is induced by σ_i , for $i = 1, \dots, n$. Then $(A^{\oplus m}, p, [\nu])$ is a Prym triple.

A symmetric correspondence. Assume that $\mathcal{T} = (A^{\oplus m}, p, [\nu])$ is a Prym triple with $p : C \rightarrow \mathbb{P}^1$ and $\nu : p^{-1}(q) \rightarrow \{1, \dots, md\}$. We shall define a symmetric correspondence on $C \times C$. Take the Galois closure $\pi : X \rightarrow \mathbb{P}^1$ of p and let H be the Galois group of the covering $X \rightarrow C$. Denote $G = \text{Gal}(\pi)$ and choose a Galois labelling $\pi^{-1}(q) \leftrightarrow G$ such that the induced labelling $p^{-1}(q) \leftrightarrow H \backslash G$ yields $\nu(H) = 1$. Define the group $\Sigma = \{\sigma_g \in S_{md} \mid g \in G\}$, where σ_g is the permutation sending $\nu(Hf) \mapsto \nu(Hfg^{-1})$ and consider the canonical homomorphism $\phi : G \rightarrow \Sigma$, $g \mapsto \sigma_g$. Since $\ker(\phi)$ is a normal subgroup of G contained in H , the minimality of π dictates that $\ker(\phi)$ is trivial, implying that ϕ is an isomorphism. Noticing that, with respect to

the Galois labelling, the monodromy group of π acts on G via multiplication on the right with the elements of G , we conclude that Σ is the monodromy group of p with respect to $[\nu]$. Hence $\phi : G \hookrightarrow \text{Aut}(A^{\oplus m})$ is a transitive representation. The fact that H is the stabilizer of the letter 1 with respect to ϕ thus implies that $\mathcal{P} = (A^{\oplus m}, r, \pi, \phi)$ for $r \in \{r_+, r_-\}$ represents Prym data. Therefore

$$D_{\mathcal{T}} := \hat{D}_p + r\Delta_C$$

is a well-defined symmetric correspondence on $C \times C$. Recall that k is the eigenvalue of the eigenvector $(1, \dots, 1)$ of A . Because $\nu(Hg) = (\phi(g))^{-1}(1)$ for all $g \in G$, equation (3.1) gives the following interpretation of $D_{\mathcal{T}}$.

Lemma 4.2. *Let $\{x_1, \dots, x_{md}\}$ be a labelling in the class $[\nu]$ and denote $A^{\oplus m} = (s_{ij})_{i,j=1}^{md}$. Then the point (x_i, x_j) appears in $D_{\mathcal{T}}$ with multiplicity s_{ij} . In particular, $D_{\mathcal{T}}$ is of bidegree (k, k) . $\square$*

In fact, if S denotes the set of non-zero entries of A and for each $s \in S$ we define a set $\hat{G}_s = \{\hat{g} \in H \backslash G / H \mid s_{(\phi(g))(1),1} = s\}$, then we find reduced divisors $D_s = \sum_{\hat{g} \in \hat{G}_s} \hat{g}$ on $C \times C$ such that $D_{\mathcal{T}} = \sum_{s \in S} s D_s$.

Definition 4.3. Let $\gamma_{\mathcal{T}} = \hat{\gamma}_p + \text{rid}_{J(C)}$. Then $P_{\pm}(\mathcal{T}) = \ker(\gamma_{\mathcal{T}} - r_{\mp} \text{id}_{J(C)})_0$ are the Prym varieties associated to $\mathcal{T}$.

Remark. Given a second labelling ν' of the fiber $p^{-1}(q)$, according to RET there exists an $f \in \text{Aut}(p)$ such that $\nu = \nu' \circ f$ if and only if ν and ν' yield the same monodromy representation for p . If such an f exists, then f induces an isomorphism of the Prym varieties.

By Proposition 3.5, if $m = 1$, then $P_+(\mathcal{T})$ and $P_-(\mathcal{T})$ are complementary subvarieties of $J(C)$ defined by $P_{\pm}(\mathcal{T}) = \text{im}(\gamma_{\mathcal{T}} - r_{\pm} \text{id}_{J(C)})$. If in addition $D_{\mathcal{T}}$ is fixed point free and $r_+ = 1$, then a theorem of Kanev (cf. [4], Theorem 3.1) states that $P_+(\mathcal{T})$ is a Prym-Tyurin variety of exponent $1 - r_-$ for C .

We now try to compute the dimension of $P_{\pm}(\mathcal{T})$. Let $\eta \in \{0, 1\}$ be such that $\eta = 1$ if $k \neq r_{\pm}$ and $\eta = 0$ else.

Proposition 4.4. *Let $\mathcal{T} = (A^{\oplus m}, p, [\nu])$ be a Prym triple with $p : C \rightarrow \mathbb{P}^1$ and assume that A has diagonal $(s, \dots, s)$. Denote $\mathcal{T}' = (A^{\oplus m} - sI_{md}, p, [\nu])$ and $\mathcal{T}_0 = (J_d^{\oplus m}, p, [\nu])$. Using the notation $d_{\pm} = \dim P_{\pm}(\mathcal{T})$ and $d_0 = \dim P_-(\mathcal{T}_0)$, we have the following identity for the dimension of $P_{\pm}(\mathcal{T})$:*

$$\pm(r_+ - r_-)d_{\pm} = (k - r_{\pm})\eta d_0 + (r_{\pm} - s)g(C) - k + s + \frac{1}{2}(D_{\mathcal{T}'} \cdot \Delta_C).$$

Proof. Denote $\gamma_0 = \gamma_{\mathcal{T}_0}$ and define an endomorphism ε on $J(C)$ such that $\varepsilon = \gamma_{\mathcal{T}} - k \text{id}_{J(C)}$ if $\eta = 1$ and $\varepsilon = \text{id}_{J(C)}$ else. Then Lemma 4.2 implies that $\varepsilon(\gamma_{\mathcal{T}} - r_+ \text{id}_{J(C)})(\gamma_{\mathcal{T}} - r_- \text{id}_{J(C)}) = 0$. Recall that $\varrho_a(\gamma_{\mathcal{T}})$ and $\varrho_a(\gamma_0)$ are self-adjoint w.r.t. the form $c_1(\Theta_C)$. Hence by direct consequence of Proposition 3.4, if $\eta = 1$ (resp. $\eta = 0$), then $\varrho_a(\gamma_{\mathcal{T}})$ has eigenvalues k, r_+, r_-

(resp. r_+, r_-) with respective multiplicities d_0, d_-, d_+ (resp. d_-, d_+). Since $\text{Tr}(\varrho_a(\gamma_{\mathcal{T}})) = \text{Tr}(\varrho_a(\gamma_{\mathcal{T}'})) + s g(C)$, we obtain

$$(4.1) \quad \begin{cases} d_+ + d_- + \eta d_0 = g(C) \\ r_- d_+ + r_+ d_- + k \eta d_0 = \text{Tr}(\varrho_a(\gamma_{\mathcal{T}'})) + s g(C) \end{cases}$$

Let $\text{Tr}_r(\gamma_{\mathcal{T}'})$ be the rational trace of $\gamma_{\mathcal{T}'}$, i.e., $\text{Tr}_r(\gamma_{\mathcal{T}'})$ is the trace of the extended rational representation $(\varrho_r \otimes 1)(\gamma_{\mathcal{T}'})$ of $H^1(C, \mathbb{Z}) \otimes \mathbb{C}$. As $\varrho_r \otimes 1$ is equivalent to $\varrho_a \oplus \overline{\varrho_a}$, it follows that $\text{Tr}_r(\gamma_{\mathcal{T}'}) = 2 \text{Tr}(\varrho_a(\gamma_{\mathcal{T}'}))$. With $D_{\mathcal{T}'}$ being of bidegree $(k-s, k-s)$, Proposition 11.5.2. of [3] implies

$$\text{Tr}(\varrho_a(\gamma_{\mathcal{T}'})) = \frac{1}{2} \text{Tr}_r(\gamma_{\mathcal{T}'}) = k - s - \frac{1}{2} (D_{\mathcal{T}'} \cdot \Delta_C).$$

Solving (4.1) for $d_{\pm}$ we obtain the desired result. $\square$

Proposition 3.5 implies that for $m = 1$ we have $d_0 = 0$. To compute $\dim P_{\pm}(\mathcal{T})$ we need to determine the intersection number $(D_{\mathcal{T}'} \cdot \Delta_C)$. We shall do this for a Prym triple $\mathcal{T} = (A^{\oplus m}, p, [\nu])$, where $A^{\oplus m} = (s_{i,j})_{i,j=1}^{md}$ has zero diagonal. Denoting the set of nonzero entries of A by S , we recall that $D_{\mathcal{T}} = \sum_{s \in S} s D_s$ with D_s reduced. Hence it suffices to determine the local intersection numbers $(D_s \cdot \Delta_C)_{(x,x)}$ at (x, x) for a ramification point $x \in C$ of $p : C \rightarrow \mathbb{P}^1$. Let $b \in \mathbb{P}^1$ be the corresponding branch point and assume that the local monodromy at b is given by $\sigma_b \in S_{md}$. Further, let $\tau \in S_{md}$ be the cycle factor of σ_b which describes the ramification at x and assume that it is of order l . For each $s \in S$ we define a set $T_{\tau,s}$ of elements $t \in \{1, \dots, l-1\}$ for which there exists a $j \in \{1, \dots, md\}$ such that $s_{j, \tau^t(j)} = s$. Then:

Lemma 4.5. *For each $s \in S$ we have $(D_s \cdot \Delta_C)_{(x,x)} = \#(T_{\tau,s})$.*

Proof. After a suitable choice of coordinates on a small open neighborhood of x the covering p is given by $z \mapsto z^l$. Then near the point (x, x) the reduced divisor D_s can be described as the union of graphs of the multiplications $z \mapsto \zeta_l^t z$ (for $t \in T_{\tau,s}$) with $\zeta_l = \exp(\frac{2\pi}{l} \sqrt{-1})$. Obviously these graphs intersect Δ_C transversally in (x, x) , thus implying $(D_s \cdot \Delta_C)_{(x,x)} = \#(T_{\tau,s})$. $\square$

Hence, given the branch locus B of p and, for each $b \in B$, the set R_b of cycle factors in the cycle decomposition of σ_b , we can calculate the intersection number as a sum

$$(D_{\mathcal{T}} \cdot \Delta_C) = \sum_{b \in B} \sum_{\tau \in R_b} \sum_{s \in S} s \#(T_{\tau,s}).$$

For an application of the lemma we refer to Example 5.3.

5. EXAMPLES

Our first example has been covered by Kanev in [5]. We will treat it by a different method.

Example 5.1. (Schläfli graph) Let $\mathcal{L}$ be the intersection graph of the 27 lines on a non-singular cubic surface in $\mathbb{P}^3$. In the notation of Example 2.1 we take τ_i ($i = 1, \dots, 5$) (resp. τ_6) to be the transformation that interchanges the rows of the double-six $M_{i,i+1}$ (resp. $M_{1,2,3}$). Denoting the adjacency matrix of $\mathcal{L}$ by A , we recall that $\text{Aut}(A) = \langle \tau_1, \dots, \tau_6 \rangle$ is a transitive subgroup of S_{27} . Note moreover that each τ_j is a reflection with exactly 15 fixed points.

Let $n \geq 7$ be an integer and choose a subset $B = \{b_1, \dots, b_{2n}\}$ of $\mathbb{P}^1$. Then we know that there exists a Prym triple $\mathcal{T} = (A, p, [\nu])$ for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and monodromy group $\text{Aut}(A)$ such that its ramification over b_i is induced by a τ_j . According to Hurwitz' formula the curve C is of genus $6n - 26$. Since no vertex v of $\mathcal{L}$ is adjacent to $\tau_j(v)$, it follows that $D_{\mathcal{T}}$ is fixed point free. As A has eigenvalues $k = 10$, $r_+ = 1$ and $r_- = -5$, Proposition 4.4 implies that $P_+(\mathcal{T})$ is an $(n - 6)$ -dimensional Prym-Tyurin variety of exponent 6 for the curve C . With regard to moduli, note that for $n = 12$ we have $g(C) = 46$, $\dim P_+(\mathcal{T}) = 6$ and $\#(B) = \dim \mathcal{A}_6 + \dim \text{Aut}(\mathbb{P}^1) = 24$, where $\mathcal{A}_6$ is the moduli space of 6-dimensional principally polarized abelian varieties.

Example 5.2. (Lattice graphs) For $n \geq 3$ let A be the adjacency matrix of the lattice graph $L_2(n)$ with vertex set $\{1, \dots, n\}^2$. We define a group $G = \langle \varphi_0, \varphi_1, \varphi_2, \varphi_3 \rangle$ generated by reflections $\varphi_h := (\tau_h, \tau_h^{-1}) \circ t$ in $\text{Aut}(A)$, where t acts on $\{1, \dots, n\}^2$ by exchange of coordinates and $\tau_0, \tau_1, \tau_2, \tau_3 \in S_n$ are given by $\tau_0 = (1)$, $\tau_1 = (1 \ n)$, $\tau_2 = (2 \ n)$ and $\tau_3 = (1 \ 2 \ \dots \ n)$. Then G is a transitive subgroup of $\text{Aut}(A)$; indeed, identifying $\{1, \dots, n\}$ and $\mathbb{Z}/n\mathbb{Z}$, we have

- a) $(\varphi_1 \circ \varphi_3)^m(1, 1) = (1, h + 1)$ for $m = 1, \dots, n - 2$;
- b) $(\varphi_3 \circ \varphi_0)(i, j) = (i + 1, j - 1)$ for $i, j = 1, \dots, n$ with $j \neq n - i$;
- c) $((\varphi_3 \circ \varphi_0)^{m-1} \circ \varphi_2)(2, 1) = (m, n - m + 1)$ for $m = 1, \dots, n$.

Given an integer $l \geq 0$, choose a subset $B = \{b_1, \dots, b_{2l+8}\}$ of $\mathbb{P}^1$. We may assume that $\mathcal{T} = (A, p, [\nu])$ is a Prym triple for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and monodromy group G such that its ramification over b_i is induced by a φ_h . Then C is of genus $(n - 1)^2 + \frac{1}{2}ln(n - 1)$. As $D_{\mathcal{T}}$ is fixed point free and A has eigenvalues $k = 2(n - 1)$, $r_+ = n - 2$ and $r_- = -2$, it follows that

$$\dim P_+(\mathcal{T}) = (n - 1)(n - 3) + \frac{1}{2}l(n - 1)(n - 2).$$

Hence, for $n = 3$ and $l \geq 1$ we obtain a finite number of finite dimensional families of l -dimensional Prym-Tyurin varieties of exponent 3 for curves of genus $3l + 4$. In anticipation of section 7 we shall say that $\mathcal{T}$ is of *type* l whenever $n = 3$ and $l \geq 1$.

Example 5.3. For an example involving symmetric correspondences with fixed points, let $n \geq 3$ and assume that A is the adjacency matrix of the graph $L_2(n)$ (the complement of $L_2(n)$) with vertex set $\{1, \dots, n\}^2$. Assume that $t \in \text{Aut}(A)$ acts on $\{1, \dots, n\}^2$ by exchange of coordinates and for

$h = 1, \dots, n-1$ define the reflection $\sigma_h := ((1 \ h+1), (1))$ in $S_n \times S_n$. We observe that $\text{Aut}(A)$ is generated by the elements t and $\sigma_1, \dots, \sigma_{n-1}$. Clearly, no vertex $(i, j) \in \{1, \dots, n\}^2$ is adjacent to $\sigma_h(i, j)$. Further, (i, j) is adjacent to $t(i, j)$ if and only if $i \neq j$.

Given nonnegative integers l_1, l_2 , we choose a subset $B = B_1 \sqcup B_2$ of $\mathbb{P}^1$ with $B_1 = \{b_{1,1}, \dots, b_{1,2(l_1+1)}\}$ and $B_2 = \{b_{2,1}, \dots, b_{2,2(l_2+n-1)}\}$. Let $\mathcal{T} = (A, p, [\nu])$ be a Prym triple for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and monodromy group $\text{Aut}(A)$ such that its ramification over $b_{1,i}$ (resp. $b_{2,j}$) is induced by t (resp. some σ_h). Then the curve C is of genus $\frac{1}{2}(n-1)(n-2) + \frac{1}{2}l_1n(n-1) + l_2n$ and Lemma 4.5 implies that $(D_{\mathcal{T}} \cdot \Delta_C) = (l_1+1)(n-1)n$. It follows that $P_+(\mathcal{T})$ is of dimension $l_1(n-1) + l_2$.

In view of moduli, note that for $l_1 = 0$ and $n \geq \frac{1}{4}(l_2^2 - 3l_2 + 6)$ we have $\dim P_+(\mathcal{T}) = l_2$ and $\#B \geq \dim \mathcal{A}_{l_2} + \dim \text{Aut}(\mathbb{P}^1)$. In particular, if $l_1 = 0$ and $n = l_2 = 6$, then $g(C) = 46$. Moreover, since $S_2 \rtimes (S_n \times S_n)$ has no subgroup of index n , Galois theory implies that no factorization $p : C \xrightarrow{n:1} C' \xrightarrow{n:1} \mathbb{P}^1$ exists.

Example 5.4. (Latin square graphs) Given an integer $n \geq 3$, we assume that A is the adjacency matrix of the Latin square graph $L_3(n)$. We recall from Example 2.5 that $(\mathbb{Z}/n\mathbb{Z})^2$ induces a transitive subgroup of $\text{Aut}(A)$ via translation; as such it coincides with $\langle (1, 1), (1, 2) \rangle$. Viewed as permutations of the vertex set $(\mathbb{Z}/n\mathbb{Z})^2$, the translations $(1, 1)$ and $(1, 2)$ split into n mutually disjoint n -cycles. For $n \geq 4$ the vertices $(i+1, j+1)$ and $(i+1, j+2)$ of $L_3(n)$ are non-adjacent to (i, j) .

Now assume that $n \geq 4$. We choose an integer $l \geq 2$ and a subset $B = \{b_1, \dots, b_{ln}\}$ of $\mathbb{P}^1$. Then there exists a Prym triple $\mathcal{T} = (A, p, [\nu])$ for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and monodromy group $(\mathbb{Z}/n\mathbb{Z})^2$ such that its ramification over b_i is induced by $(1, 1)$ or $(1, 2)$. We find that C is of genus $1 - n^2 + \frac{1}{2}ln^2(n-1)$. Moreover, since $\deg(p) = \#(\mathbb{Z}/n\mathbb{Z})^2$, it is immediately seen that p is a Galois covering. Using the fact that $D_{\mathcal{T}}$ is fixed point free and A has eigenvalues $k = 3(n-1)$, $r_+ = n-3$ and $r_- = -3$, we compute

$$\dim P_+(\mathcal{T}) = -(n-1)(n-2) + \frac{1}{2}ln(n-1)(n-3).$$

Hence, for $n = 4$ we get finitely many finite dimensional families of $6(l-1)$ -dimensional Prym-Tyurin varieties of exponent 4 for curves of genus $24l-15$.

6. A SPLITTING

We show that for certain Prym triples $\mathcal{T} = (A^{\oplus m}, p, [\nu])$ the covering $p : C \rightarrow \mathbb{P}^1$ splits into a covering $f : C \rightarrow C'$ of degree d and a covering $h : C' \rightarrow \mathbb{P}^1$ of degree m such that f depends essentially on $D_{\mathcal{T}}$. Recall that k is the eigenvalue of the eigenvector $(1, \dots, 1)$ of A .

Theorem 6.1. *Assume that $A \in \{0, 1\}^{d \times d}$ is a Prym matrix with zero diagonal and eigenvalue k of multiplicity 1. Given $m \geq 2$, let $\mathcal{T} = (A^{\oplus m}, p, [\nu])$*

be a Prym triple for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B . Denote $C_0 := p^{-1}(\mathbb{P}^1 \setminus B)$. Then there exists a unique splitting

$$p : C \xrightarrow[f]{d:1} C' \xrightarrow[h]{m:1} \mathbb{P}^1$$

such that, for any $(x, x') \in (C_0 \times C_0) \setminus \Delta_{C_0}$, the points x, x' are in the same fiber of f if and only if there is a finite sequence of points $x = x_0, \dots, x_l = x'$ on C_0 with $(x_j, x_{j+1}) \in D_{\mathcal{T}}$ for all $j = 0, \dots, l-1$.

Proof. Fix a point $q_0 \in \mathbb{P}^1 \setminus B$ and assume that ν is a labelling of the fiber $p^{-1}(q_0)$. We denote $S = \{1, \dots, m\}$, $T = \{1, \dots, d\}$ and identify $S \times T$ with $\{1, \dots, md\}$ via the bijection $(s, t) \leftrightarrow (s-1)m + t$. Then ν turns into a bijection $(\nu_1, \nu_2) : p^{-1}(q_0) \rightarrow S \times T$. Let Σ be the monodromy group of p with respect to (ν_1, ν_2) and split its elements accordingly into $\sigma = (\sigma_1, \sigma_2)$. Denoting $A := (a_{i,j})_{i,j=1}^d$, we may view $A^{\oplus m}$ as the matrix of entries $c_{u,u'}$, where $u = (s, t)$ and $u' = (s', t')$ run through the set $S \times T$, such that $c_{u,u'} = a_{t,t'}$ if $s = s'$ and $c_{u,u'} = 0$ else. According to Proposition 2.3 the matrix A is the adjacency matrix of a connected strongly regular graph $\mathcal{G}$ on d vertices. Thus, for $u = (s, t)$ and $u' = (s', t')$ there exists a finite sequence $u = u_0, \dots, u_l = u'$ in $S \times T$ such that $c_{u_j, u_{j+1}} = 1$ for all $j = 0, \dots, l-1$ if and only if $s = s'$. Hence $\sigma_1(\cdot, t) = \sigma_1(\cdot, t')$ for all $\sigma \in \Sigma$, i.e., there is a unique $\tau_\sigma \in S_m$ such that $\sigma_1(\cdot, t) = \tau_\sigma(\cdot)$ for all $t \in T$.

Let $\pi : X \rightarrow \mathbb{P}^1$ be the Galois closure of p and denote $G = \text{Gal}(\pi)$. As we have seen in section 4, there exists an isomorphism $\phi : G \rightarrow \Sigma$ such that the Galois group H of $X \rightarrow C$ is the stabilizer of $(1, 1) \in S \times T$ w.r.t. ϕ and any Galois labelling of a fiber of π induces a labelling in the class $[\nu]$ via the identification $Hg \leftrightarrow g^{-1}(1, 1)$. We let H' be the stabilizer of $1 \in S$ with respect to $\psi \circ \phi$, where $\psi : \Sigma \rightarrow S_m$ is the transitive representation induced by $\sigma \mapsto \tau_\sigma$. Write $C' = X/H'$; since $H \subset H'$ (resp. $H' \subset G$) is a subgroup of index d (resp. m), there are canonical coverings $f : C \rightarrow C'$ of degree d and $h : C' \rightarrow \mathbb{P}^1$ of degree m such that $p = h \circ f$. Take a point $q \in \mathbb{P}^1 \setminus B$ and a Galois labelling $\pi^{-1}(q) \leftrightarrow G$. For any element $g \in G$, if $(s, t) = g^{-1}(1, 1)$, then $H'g \leftrightarrow g^{-1}(1) = s$, i.e., on the fiber $p^{-1}(q)$ the covering f is given by $(s, t) \mapsto s$. With reference to Lemma 4.2 we conclude that f has the desired properties. Using the monodromy of p , the reader will easily check that the splitting is unique. $\square$

With $\mathcal{T}$, f and h as above, we say that the pair of coverings (f, h) represents the *canonical splitting* for $\mathcal{T}$.

Corollary 6.2. *For integers $d, m \geq 2$, assume that $\mathcal{T} = ((J_d - I_d)^{\oplus m}, p, [\nu])$ is a Prym triple associated to a covering $p : C \rightarrow \mathbb{P}^1$ and let (f, h) be its canonical splitting. Then $P_+(T)$ is the usual Prym variety associated to the covering f , i.e., $P_+(T)$ and $\text{im} f^*$ are complementary subvarieties of $J(C)$.*

Proof. According to Theorem 6.1 we have $D_T(x) = -x + f^*f(x)$ for all $x \in C$ in an unramified fiber of p . Hence $\gamma_T + \text{id}_{J(C)} = f^*N_f$ and thus $P_-(T) = \text{im } f^*$. As $(J_d - I_d)^{\oplus m}$ is a Prym matrix, Proposition 3.5 implies that $P_+(T)$ and $\text{im } f^*$ are complementary in $J(C)$. $\square$

Corollary 6.2 has a natural converse. Before addressing this, we recall that a smooth projective curve of genus g is m -gonal for all $m \geq \lceil \frac{g}{2} \rceil + 1$ (cf. [1], Existence Theorem, p. 206).

Corollary 6.3. *Assume that $f : C \rightarrow C'$ is a covering of degree $d \geq 2$ of a curve C' of genus $g \geq 1$ and let $h : C' \rightarrow \mathbb{P}^1$ be a covering of degree $m \geq \lceil \frac{g}{2} \rceil + 1$. Then there exists a labelling class $[\nu]$ for the covering $h \circ f$ such that $\mathcal{T} = ((J_d - I_d)^{\oplus m}, h \circ f, [\nu])$ is a Prym triple and $P_+(\mathcal{T})$ is the usual Prym variety associated to f .*

Proof. Take a point $q \in \mathbb{P}^1$ outside the branch locus of $h \circ f$. Then we can define a bijection $\nu = (\nu_1, \nu_2) : (h \circ f)^{-1}(q) \rightarrow \{1, \dots, m\} \times \{1, \dots, d\}$ such that $\nu_1(x) = \nu_1(x') \iff f(x) = f(x')$, for all $x, x' \in (h \circ f)^{-1}(q)$. It is immediately seen that $((J_d - I_d)^{\oplus m}, h \circ f, [\nu])$ represents a Prym triple with canonical splitting (f, h) . Now apply Corollary 6.2. $\square$

Given integers $d, m \geq 2$, assume that $\mathcal{T} = ((J_d - I_d)^{\oplus m}, p, [\nu])$ is a Prym triple. We shall call $\mathcal{T}$ *simple* if its canonical splitting (f, h) is simple, i.e., if f and h are simply branched coverings such that no ramified fiber of h contains a branch point of f and no unramified fiber of h contains more than one branch point of f . It should be noted that simplicity can also be described in terms of the monodromy of p alone, without reference to f and h .

To conclude this section, we use simple Prym triples to characterize (at least up to isogeny) abelian varieties corresponding to the general popints of $\mathcal{A}_4$ and $\mathcal{A}_5$.

Lemma 6.4. (1) *The general 4-dimensional principally polarized abelian variety is isogenous to a Prym variety $P_+(\mathcal{T})$ for a simple Prym triple $\mathcal{T} = ((J_2 - I_2)^{\oplus 3}, p, [\nu])$ such that the covering p has exactly 4 simple and 10 double branch points.*

(2) *The general 5-dimensional principally polarized abelian variety is of the form $P_+(\mathcal{T})$, where $\mathcal{T} = ((J_2 - I_2)^{\oplus 4}, p, [\nu])$ is a simple Prym triple such that the covering p has exactly 18 double branch points.*

Proof. For integers $g \geq 1$ and $n \geq 0$, let $\mathcal{R}(g, n)$ be the moduli space of equivalence classes of double coverings $f : C \rightarrow C'$ with C' of genus g and f branched at $2n$ distinct points of C' . We shall need the following fact: Let m be an integer. If $m \geq \lceil \frac{g}{2} \rceil + 1$, then for a double covering $f : C \rightarrow C'$ corresponding to a general point of $\mathcal{R}(g, n)$ there exists an m -fold covering $h : C' \rightarrow \mathbb{P}^1$ such that the covering pair (f, h) is simple. The proof is left

to the reader. As in [2], p. 122, we let $p_{(g,n)} : \mathcal{R}(g,n) \rightarrow \mathcal{A}_{g+n-1}(\delta)$ be the usual Prym morphism, where $\mathcal{A}_{g+n-1}(\delta)$ is the moduli space of abelian g -folds with polarization type δ . According to [2], Theorem 2.2, the morphism $p_{(3,2)} : \mathcal{R}(3,2) \rightarrow \mathcal{A}_4(1,2,2,2)$ is dominant. Moreover, for the general double covering $f : C \rightarrow C'$ with 4 branch points and $g(C') = 3$ there exists a 3-fold covering $h : C' \rightarrow \mathbb{P}^1$ such that the pair (f,h) is simple and the covering $h \circ f$ has exactly 4 simple and 10 double branch points. Together with Corollary 6.3 this shows (1). To prove (2) we recall that $p_{(6,0)} : \mathcal{R}(6,0) \rightarrow \mathcal{A}_5$ is dominant (cf. [10]). Hence it suffices to note that for the general étale double covering $f : C \rightarrow C'$ with $g(C') = 6$ there exists a 4-fold covering $h : C' \rightarrow \mathbb{P}^1$ branched at 18 points such that the pair (f,h) is simple. $\square$

7. PRYM-TYURIN VARIETIES OF ARBITRARY EXPONENT ≥ 3

We show how the graph $\overline{L_2(n)} \in \text{SRG}(n^2, (n-1)^2, (n-2)^2, (n-1)(n-2))$ for $n \geq 3$ can be employed to construct families of Prym-Tyurin varieties of exponent n . These varieties turn out to be the product of the Jacobians of two n -gonal curves.

Example 7.1. Given an integer $n \geq 3$, we shall try to construct Prym-Tyurin varieties of exponent n . Assume that A is the adjacency matrix of the graph $\overline{L_2(n)}$ with vertex set $\{1, \dots, n\}^2$. Recall that $S_n \times S_n$ is a transitive subgroup of $\text{Aut}(A)$. For $i = 1, \dots, n-1$ we define reflections $\sigma_{1,i} := ((1 \ i+1), (1))$ and $\sigma_{2,i} := ((1), (1 \ i+1))$ in $S_n \times S_n$. Note that $S_n \times S_n$ is freely generated by these reflections.

Given nonnegative integers l_1, l_2 such that $l_1 + l_2 \geq 1$, let $B = B_1 \sqcup B_2$ be a finite subset of $\mathbb{P}^1$ with $B_m = \{b_{m,1}, \dots, b_{m,2(l_m+n-1)}\}$. Assume that $\mathcal{T} = (A, p, [\nu])$ is a Prym triple for a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and monodromy group $S_n \times S_n$ such that its ramification over $b_{m,i}$ is induced by a $\sigma_{m,h}$; in this situation we say that $\mathcal{T}$ is of *type* (l_1, l_2) . Since no vertex (i, j) of $\overline{L_2(n)}$ is adjacent to $\sigma_{m,h}(i, j)$, the correspondence $D_{\mathcal{T}}$ is fixed point free. Moreover, it is easily seen that all $\sigma_{m,h}$ decompose into n mutually disjoint transpositions on the set $\{1, \dots, n\}^2$. Hence, as A has eigenvalues $k = (n-1)^2$, $r_+ = 1$ and $r_- = -n+1$, it follows that $P_+(\mathcal{T})$ is an $(l_1 + l_2)$ -dimensional Prym-Tyurin variety of exponent n for the curve C of genus $(n-1)^2 + (l_1 + l_2)n$.

Recall from Example 5.2 that a Prym triple $\mathcal{T}$ of type l yields an l -dimensional Prym-Tyurin variety $P_+(\mathcal{T})$ of exponent 3. We will show that for $n = 3$ the Prym-Tyurin varieties of the preceding example are the same as those of Example 5.2. More precisely, let A be the adjacency matrix of the lattice graph $L_2(3)$ and assume that $[\nu]$ is a labelling class for a covering $p : C \rightarrow \mathbb{P}^1$ of degree 9. Given the isomorphism of graphs $\xi : L_2(3) \xrightarrow{\sim} \overline{L_2(3)}$ induced by the matrix $\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ as in Example 2.4, we have:

Lemma 7.2. *Let $l \geq 1$ be an integer. Then $\mathcal{T} = (A, p, [\nu])$ is a Prym triple of type l if and only if there exist integers $l_1, l_2 \geq 0$ such that $l_1 + l_2 = l$ and*

$\mathcal{T}' = (J_9 - I_9 - A, p, [\xi^{-1} \circ \nu])$ is a Prym triple of type (l_1, l_2) . In particular, if $\mathcal{T}$ is of type l , then $P_{\pm}(\mathcal{T}) = P_{\pm}(\mathcal{T}')$.

Proof. It suffices to note that the reflections $\varphi_0, \varphi_1, \varphi_2, \varphi_3 \in \text{Aut}(L_2(3))$ of Example 5.2 satisfy the identities $\xi^{-1}\sigma_{1,1}\xi = \varphi_0$, $\xi^{-1}\sigma_{1,2}\xi = \varphi_0\varphi_3\varphi_0^{-1}$, $\xi^{-1}\sigma_{2,1}\xi = \varphi_1\varphi_2\varphi_1^{-1}$ and $\xi^{-1}\sigma_{2,2}\xi = \varphi_2$. $\square$

For Prym-Tyurin varieties $P_{+}(\mathcal{T})$ with $\mathcal{T}$ of type (l_1, l_2) we have the following characterization.

Theorem 7.3. *Assume that A is the adjacency matrix of $\overline{L_2(n)}$ with $n \geq 3$. For nonnegative integers l_1, l_2 such that $l_1 + l_2 \geq 1$, let $\mathcal{T}$ be a Prym triple of type (l_1, l_2) associated to A and a covering $C \rightarrow \mathbb{P}^1$. Then there exist n -gonal curves C_1 of genus l_1 and C_2 of genus l_2 such that $C = C_1 \times_{\mathbb{P}^1} C_2$ and $P_{+}(\mathcal{T}) \simeq J(C_1) \times J(C_2)$.*

Proof. Let $p : C \rightarrow \mathbb{P}^1$ and $[\nu]$ be the covering and labelling class such that $\mathcal{T} = (A, p, [\nu])$. Then $S_n \times S_n$ is the monodromy group of p . We take the inclusion $\iota : S_n \times S_n \hookrightarrow \text{Perm}(N \times N)$ and write $N = \{1, \dots, n\}$. Let $\pi : X \rightarrow \mathbb{P}^1$ be the Galois closure of p and denote $G = \text{Gal}(\pi)$. Then there exists an isomorphism $\phi : G \rightarrow \Sigma$ such that the Galois group H of $X \rightarrow C$ is the stabilizer of $(1, 1) \in N \times N$ w.r.t. $\iota \circ \phi$ and any Galois labelling of a fiber of π induces a labelling in the class $[\nu]$ via the identification $Hg \leftrightarrow g^{-1}(1, 1)$. Take the projection mappings $\text{pr}_1, \text{pr}_2 : S_n \times S_n \rightarrow S_n$ onto the first and second factor and let H_1 (resp. H_2) be the stabilizer of the letter $1 \in N$ w.r.t. $\phi_1 := \text{pr}_1 \circ \phi$ (resp. $\phi_2 := \text{pr}_2 \circ \phi$). Observing that $H = H_1 \cap H_2$, we take the quotient curves $C_m = X/H_m$ for $m = 1, 2$ and let $f_m : C \rightarrow C_m$ and $h_m : C_m \rightarrow \mathbb{P}^1$ be the canonical coverings. The transitivity of ϕ_m implies that f_m and h_m are of degree n . In addition to $H = H_1 \cap H_2$ we have $G = \langle H_1, H_2 \rangle$. Using elementary Galois theory we thus find

$$\mathbb{C}(C) = \mathbb{C}(C_1) \otimes_{\mathbb{C}(\mathbb{P}^1)} \mathbb{C}(C_2).$$

Hence C is the fiber product of the n -gonal curves C_1 and C_2 with projection morphisms f_1 and f_2 . Let B be the branch locus of the covering p . Then h_m for $m = 1, 2$ is a simple covering with branch locus B_m , where B_m is the set of points $b \in B$ such that, in the notation of Example 7.1, the local monodromy of p is given by a permutation $\sigma_{m,i}$. As $\#(B_m) = 2(l_m + n - 1)$, we find $g(C_m) = l_m$.

It remains to show that $P_{+}(\mathcal{T}) \simeq J(C_1) \times J(C_2)$. Choose a point $q \in \mathbb{P}^1 \setminus B$ and a Galois labelling $\pi^{-1}(q) \leftrightarrow G$. Then take a labelling $\{y_{1,1}, \dots, y_{n,n}\}$ for $p^{-1}(q)$ and a labelling $\{z_{m,1}, \dots, z_{m,n}\}$ for $h_m^{-1}(q)$, $m = 1, 2$ such that $y_{g^{-1}(1,1)} \leftrightarrow Hg$ and $z_{m,(\phi_m(g))^{-1}(1)} \leftrightarrow H_m g$, for all $g \in G$. We observe that $f_1^{-1}(z_{1,s}) = \{y_{s,j} \mid j \in N\}$ and $f_2^{-1}(z_{2,t}) = \{y_{i,t} \mid i \in N\}$, for all $s, t \in N$.

According to Lemma 4.2 we have $D_T(y_{s,t}) = \sum_{i \neq s, j \neq t} y_{i,j}$ and therefore

$$f_1^* z_{1,s} + f_2^* z_{2,t} = \sum_{j \in N} y_{s,j} + \sum_{i \in N} y_{i,t} = p^* p(y_{s,t}) + y_{s,t} - D_T(y_{s,t}).$$

Hence, for $y, y' \in p^{-1}(\mathbb{P}^1 \setminus B)$ and $z_m = f_m(y)$, $z'_m = f_m(y')$ with $m = 1, 2$ we obtain, using divisor class notation,

$$f_1^*[z_1 - z'_1] + f_2^*[z_2 - z'_2] = -(\gamma_T - \text{id}_{J(C)})([y - y']).$$

Consequently, defining $\varphi = f_1^* \psi_1 + f_2^* \psi_2 : J(C_1) \times J(C_2) \rightarrow J(C)$, where $\psi_m : J(C_1) \times J(C_2) \rightarrow J(C_m)$ is the projection on the m -th factor, we get $P_+(T) = \text{im}(\gamma_T - \text{id}_{J(C)}) \subset \text{im} \varphi$. Because $\dim P_+(T) = l_1 + l_2 = \dim J(C_1) + \dim J(C_2)$, it thus follows that $\varphi : J(C_1) \times J(C_2) \rightarrow P_+(T)$ is an isogeny. As $P_+(T)$ is a Prym-Tyurin variety of exponent n for C , the restriction of the canonical polarization Θ_C to $P_+(T)$ is of type $(n, \dots, n)$. Lemma 12.3.1 of [3] implies that the polarization $\varphi^* \Theta_C$ of $J(C_1) \times J(C_2)$ is of type $(n, \dots, n)$, as well. Hence $\varphi : J(C_1) \times J(C_2) \rightarrow P_+(T)$ is an isogeny of degree 1, i.e., an isomorphism. $\square$

Remark. In spite of the similarities between the Examples 5.4 and 7.1, the preceding theorem does not fully extend to Prym triples $\mathcal{T}$ such as in Example 5.4. In fact, defining n -gonal curves C_1 and C_2 analogously to those in the proof, we get $C = C_1 \times_{\mathbb{P}^1} C_2$. A simple computation shows, however, that the dimensions of $P_+(T)$ and $J(C_1) \times J(C_2)$ do not match.

A different construction. In [6], Lange, Recillas and Rochas define non-trivial families of Prym-Tyurin varieties of exponent 3. Here is a recap of their construction: Given a hyperelliptic curve X of genus $g \geq 3$ and an étale covering $f : \tilde{X} \rightarrow X$ of degree 3, let $h : X \rightarrow \mathbb{P}^1$ be the covering given by the g_2^1 and define the curve $C = (f^{(2)})^{-1}(g_2^1)$, where $f^{(2)} : \tilde{X}^{(2)} \rightarrow X^{(2)}$ is the second symmetric product of f . Assume for the moment that C is smooth and irreducible. Denote $\tilde{C} = \mu^{-1}(C)$, where $\mu : \tilde{X}^2 \rightarrow \tilde{X}^{(2)}$ is the canonical $2 : 1$ map and let $\pi : \tilde{C} \rightarrow \tilde{X}$ be the projection on the first factor, where $\tilde{C}$ is considered as a curve in $\tilde{X}^2$. Now define the covering $p : C \xrightarrow{9:1} \mathbb{P}^1$ induced by $h \circ f \circ \pi : \tilde{C} \rightarrow \mathbb{P}^1$ and let $\iota : X \rightarrow X$ be the hyperelliptic involution. Then p , $h \circ f$ and h have the same branch locus B , which may be assumed to be of cardinality $2l + 8$ for $l \geq 0$. To obtain a divisorial correspondence on $C \times C$, we choose a point $q \in \mathbb{P}^1 \setminus B$ and denote the fiber $h^{-1}(q)$ by $\{x, \iota(x)\}$. Write $f^{-1}(x) = \{y_1, y_2, y_3\}$ and $f^{-1}(\iota(x)) = \{z_1, z_2, z_3\}$; then $p^{-1}(q) = \{y_i + z_j \mid i, j = 1, 2, 3\}$ and the identity

$$D(y_s + z_t) = \sum_{j \neq t} (y_s + z_j) + \sum_{i \neq s} (y_i + z_t)$$

defines a fixed point free symmetric correspondence D of bidegree $(2, 2)$ on $C \times C$. Note that the matrix of entries $a_{(s,j),(i,t)}$ (for $(s, j), (i, t) \in \{1, 2, 3\}^2$)

given by

$$a_{(s,j),(i,t)} = \begin{cases} 1 & \text{if } (y_s + z_j, y_i + z_t) \in D \\ 0 & \text{else} \end{cases}$$

is the adjacency matrix of $L_2(3)$. Hence the canonical endomorphism γ_D of $J(C)$ satisfies the equation

$$(\gamma_D - \text{id}_{J(X)})(\gamma_D + 2 \text{id}_{J(X)}) = 0.$$

For $l \geq 1$ it follows that $P := \text{im}(\gamma_D - \text{id}_{J(C)})$ is an l -dimensional Prym-Tyurin variety of exponent 3 for the curve C of genus $3l + 4$. We shall call P (resp. p) the Prym variety (resp. covering) associated to f and h .

The preceding construction is a special case of Example 7.1. This is a direct consequence of Lemma 7.2 and the following result, for integers $l \geq 1$.

Proposition 7.4. *The Lange-Recillas-Rochas family of l -dimensional Prym-Tyurin varieties of exponent 3 coincides with the family of Prym-Tyurin varieties $P_+(\mathcal{T})$ for Prym triples $\mathcal{T}$ of type l .*

Proof. Let P (resp. $p : C \rightarrow \mathbb{P}^1$) be the Prym variety (resp. covering) associated to an étale threefold covering $f : \tilde{X} \rightarrow X$ and a double covering $h : X \rightarrow \mathbb{P}^1$ with branch locus B of cardinality $2l + 8$. We fix a point $q \in \mathbb{P}^1 \setminus B$ and write $N = \{1, 2, 3\}$. Using the notation of the preceding construction, we define the bijections $\nu : p^{-1}(q) \rightarrow N \times N$, $y_i + z_j \mapsto (i, j)$ and $\mu : (h \circ f)^{-1}(q) \rightarrow \{1, 2\} \times N$, sending $y_i \mapsto (1, i)$ and $z_j \mapsto (2, j)$. We let ρ (resp. ϱ) be the monodromy representations for p (resp. $h \circ f$) induced by ν (resp. μ). Choose a small q -based loop $\lambda \subset \mathbb{P}^1 \setminus B$ around a point $b \in B$. Then $\varrho([\lambda]) = v_1 v_2 v_3$ is the product of mutually disjoint transpositions $v_j = ((1, s_j) (2, t_j))$, $s_j, t_j \in N$. Employing the fact that p comes from $h \circ f \circ \pi$ with π as in the construction, one easily checks that $\xi \rho([\lambda]) \xi^{-1}$ is a conjugate of some $\sigma_{m,h} \in S_3 \times S_3$ (in the notation of Example 7.1). By transitivity of $\text{im } \rho$ it thus follows that $\xi(\text{im } \rho) \xi^{-1} = S_3 \times S_3$. Hence, if A denotes the adjacency matrix of the graph $L_2(3)$, then $\mathcal{T} = (A, p, [\nu])$ is a Prym triple of type l and Lemma 4.2 implies that $P = P_+(\mathcal{T})$.

Conversely, let $\mathcal{T}$ be a Prym triple of type l associated to a covering $p : C \rightarrow \mathbb{P}^1$ with branch locus B and a labelling class $[\nu]$, where $\nu : p^{-1}(q) \rightarrow \{1, 2, 3\}^2$ is a labelling for the fiber of p over a point $q \in \mathbb{P}^1 \setminus B$. Given the monodromy representation $\rho : \pi_1(\mathbb{P}^1 \setminus B, q) \rightarrow \text{Perm}(\{1, 2, 3\}^2)$ for p induced by ν , we take coordinate mappings ρ_1, ρ_2 such that $\rho(\beta)$ splits as $(\rho_1(\beta), \rho_2(\beta))$ for $\beta \in \pi_1(\mathbb{P}^1 \setminus B, q)$. We then have a transitive representation $\varrho : \pi_1(\mathbb{P}^1 \setminus B, q) \rightarrow \text{Perm}(\{1, 2\} \times \{1, 2, 3\})$, defined by the relations $\varrho(\beta)(1, i) = (2, \rho_2(\beta)(i, 1))$ and $\varrho(\beta)(2, j) = (1, \rho_1(\beta)(1, j))$. Using the local monodromy of p , one shows by analogy with the proof of Theorem 6.1 that ϱ is a monodromy representation for a covering

$$h \circ f : \tilde{X} \xrightarrow{3:1} X \xrightarrow{2:1} \mathbb{P}^1,$$

where f is étale and $g(X) \geq 4$. Then ρ is immediately seen to act as a monodromy representation for the covering that is associated to f and h . Hence $P_+(\mathcal{T})$ (resp. p) is the Prym variety (resp. covering) associated to f and h . $\square$

REFERENCES

- [1] E. Arbarello, M. Cornalba, P.A. Griffiths, J. Harris, *Geometry of Algebraic Curves; Volume I*, Springer-Verlag, 1985
- [2] F. Bardelli, C. Ciliberto, A. Verra: *Curves of minimal genus on a general abelian variety*, *Comp. Math.* **96** (1995), 115–147.
- [3] C. Birkenhake, H. Lange, *Complex Abelian Varieties*, Springer-Verlag, 2004.
- [4] V. Kanev: *Principal polarizations of Prym-Tjurin varieties*, *Comp. Math.* **64** (1987), 243–270.
- [5] V. Kanev: *Spectral curves and Prym-Tjurin varieties I*, *Proceedings of the Egloffstein conference 1993*, de Gruyter (1995), 151–198.
- [6] H. Lange, S. Recillas, A.M. Rochas: *A family of Prym-Tyurin varieties of exponent 3*, [arXiv:math.AG/0412103](https://arxiv.org/abs/math/0412103).
- [7] Y.I. Manin, *Cubic forms: Algebra, Geometry, Arithmetic*, North-Holland, 1974.
- [8] J.Y. Mérindol: *Variétés de Prym d'un revêtement galoisien*, *J. reine angew. Math.* **461** (1995), 49–61.
- [9] R. Miranda, *Algebraic Curves and Riemann Surfaces*, American Mathematical Society, 1995.
- [10] D. Mumford: *Prym varieties I*, in: *Contributions to Analysis*, Academic Press (1974), 325–350.
- [11] J.H. van Lint, R.M. Wilson, *A course in Combinatorics*, Cambridge University Press, 2002.
- [12] H. Völklein, *Groups as Galois groups*, Cambridge University Press, 1996.

FACULTEIT WISKUNDE EN INFORMATICA, UNIVERSITY OF AMSTERDAM, PLANTAGE
MUIDERGRACHT 24, 1018 TV AMSTERDAM, THE NETHERLANDS.

E-mail address: `rsalomon@science.uva.nl`